



March 28, 2022

U.S. Department of Labor  
Occupational Safety and Health Administration  
200 Constitution Avenue, NW  
Washington, D.C. 20210

**Re: Docket No. OSHA-2020-0004**

To Whom It May Concern:

On behalf of the American Association of Oral and Maxillofacial Surgeons (AAOMS), the professional association that represents more than 9,000 oral and maxillofacial surgeons (OMSs) in the United States, we are pleased to provide comment on the Occupational Safety and Health Administration's Emergency Temporary Standard in response to the March 23, 2022, Federal Register notice. We commend OSHA for its efforts throughout the pandemic to ensure safe conditions for our nation's workers and the particular concern the agency has shown for our nation's healthcare personnel.

OMSs are surgically and medically trained dental specialists who treat conditions, defects, injuries and esthetic aspects of the mouth, teeth, jaws, neck and face. After earning a dental degree from an accredited four-year dental school, OMSs complete a minimum of four years of hospital-based oral and maxillofacial surgery residency training, which includes rotations in such areas as general surgery, anesthesia and clinical research. As the surgical branch of dentistry, our members provided frontline response to address the needs of emergent dental patients during the pandemic.

Throughout the pandemic, our goal has been to ensure that dental patients are treated outside of the emergency room so our nation's hospitals are free to respond to pandemic patients. These dental emergencies – which can be life threatening if left untreated – include treatment of facial trauma, dental pain, abscesses and other pathology, as well as dental treatment required prior to critical medical procedures. To treat these conditions, OMSs must work in close proximity to patients and perform aerosol-generating procedures.

Early in the pandemic, the uncertainty surrounding how SARS-CoV-2 spread forced most dental practices – including OMS offices – to temporarily close. As we all learned more about the virus, we were able to implement protocols that would protect not only our patients but our healthcare staff. This allowed us to transition from the use of respirators – such as N95 masks – when treating all patients, to strategic use based on patient vaccination status, clinical pre-operative testing and community transmission rates.

A recent study from the Ohio State University ([Attachment 1](#)) supported this recommendation, concluding that “when infection control measures are used, such as preoperative mouth rinses and intraoral high-volume evacuation, dental treatment is not a factor in increasing the risk for transmission of SARS-CoV-2 in asymptomatic patients and that standard infection control practices are sufficiently capable of protecting personnel and patients from exposure to potential pathogens.”<sup>1</sup> Due to this and other similar studies that indicate dental aerosolizing procedures are not at risk, AAOMS believes that rather than utilizing respirators, the use of fluid resistant masks or mask covers, eye protection and face shields provides adequate protection for the treatment of asymptomatic patients in the dental and OMS office. AAOMS urges OSHA to consider this study’s findings and our position when finalizing the emergency temporary standard and transitioning it to a permanent regulation.

On behalf of AAOMS, thank you for your consideration. For questions or additional information, please contact Sandy Guenther, manager of state government affairs and advocacy engagement at 800-822-6637, ext. 4388 or [sguenther@aaoms.org](mailto:sguenther@aaoms.org).

Sincerely,

A handwritten signature in black ink that reads "J. David Johnson Jr., DDS". The signature is written in a cursive style with a large, stylized initial "J".

J. David Johnson Jr., DDS  
AAOMS President

---

<sup>1</sup> Meethil AP, Saraswat S, Chaudhary PP, Dabdoub SM, Kumar PS. Sources of SARS-CoV-2 and Other Microorganisms in Dental Aerosols. *Journal of Dental Research*. 2021;100(8):817-823.  
doi:10.1177/00220345211015948

# Sources of SARS-CoV-2 and Other Microorganisms in Dental Aerosols

Journal of Dental Research

1–7

© International &amp; American Associations for Dental Research 2021



Article reuse guidelines:

[sagepub.com/journals-permissions](https://sagepub.com/journals-permissions)

DOI: 10.1177/00220345211015948

[journals.sagepub.com/home/jdr](https://journals.sagepub.com/home/jdr)

A.P. Meethil<sup>1</sup>, S. Saraswat<sup>1</sup>, P.P. Chaudhary<sup>2</sup>, S.M. Dabdoub<sup>1</sup> , and P.S. Kumar<sup>1,3</sup> 

## Abstract

On March 16, 2020, 198,000 dentists in the United States closed their doors to patients, fueled by concerns that aerosols generated during dental procedures are potential vehicles for transmission of respiratory pathogens through saliva. Our knowledge of these aerosol constituents is sparse and gleaned from case reports and poorly controlled studies. Therefore, we tracked the origins of microbiota in aerosols generated during ultrasonic scaling, implant osteotomy, and restorative procedures by combining reverse transcriptase quantitative polymerase chain reaction (to identify and quantify SARS-CoV-2) and 16S sequencing (to characterize the entire microbiome) with fine-scale enumeration and source tracking. Linear discriminant analysis of Bray-Curtis dissimilarity distances revealed significant class separation between the salivary microbiome and aerosol microbiota deposited on the operator, patient, assistant, or the environment ( $P < 0.01$ , analysis of similarities). We also discovered that 78% of the microbiota in condensate could be traced to the dental irrigant, while saliva contributed to a median of 0% of aerosol microbiota. We also identified low copy numbers of SARS-CoV-2 virus in the saliva of several asymptomatic patients but none in aerosols generated from these patients. Together, the bacterial and viral data encourage us to conclude that when infection control measures are used, such as preoperative mouth rinses and intraoral high-volume evacuation, dental treatment is not a factor in increasing the risk for transmission of SARS-CoV-2 in asymptomatic patients and that standard infection control practices are sufficiently capable of protecting personnel and patients from exposure to potential pathogens. This information is of immediate urgency, not only for safe resumption of dental treatment during the ongoing COVID-19 pandemic, but also to inform evidence-based selection of personal protection equipment and infection control practices at a time when resources are stretched and personal protection equipment needs to be prioritized.

**Keywords:** ultrasonics, dental implants, irrigant, saliva, microbiota, DNA sequence analysis

## Introduction

On March 11, 2020, the World Health Organization declared coronavirus disease 2019 (COVID-19) a pandemic. On March 16, the American Dental Association responded with a recommendation that dental offices refrain from providing nonemergency services, resulting in 198,000 dentists closing their doors to patients in the United States alone. Nine months later, restrictions on the use of key instruments and procedures are still in place, fueling concern in providers and patients alike (Zhao et al. 2020). This is despite the fact that 3.5 billion individuals experience caries, periodontitis, or oral cancer (Kassebaum et al. 2014)—diseases that not only impede essential functions, such as speech and mastication (National Institute of Dental and Craniofacial Research 2014), but can also influence the course and outcomes of diabetes, atherosclerosis, and rheumatoid arthritis (D’Aiuto et al. 2005; de Pablo et al. 2009; Tonetti et al. 2013). Indeed, emerging evidence suggests that untreated periodontitis might increase the risk for developing severe forms of COVID-19 (Gupta and Sahni 2020; Pitones-Rubio et al. 2020).

Regulators and health authorities based this guidance largely on literature that aerosols generated during medical procedures, such as intubation/extubation, bronchoscopy, ventilation, and airway suctioning, can transmit infections (Davies

et al. 2009; Judson and Munster 2019; Centers for Disease Control and Prevention 2020). Although microorganisms have been identified in dental aerosols (Grenier 1995; Harrel et al. 1998), our recent review found no information regarding their origins (Kumar and Subramanian 2020). Therefore, there is an urgent need to inform infection control science by definitively identifying the source of bacteria and viruses in aerosol-generating dental procedures (AGDPs).

We combined a rigorous clinical study with microbiomics and robust ecologic statistics to test the hypothesis that microorganisms from AGDPs are of nonsalivary origin. This

<sup>1</sup>Division of Periodontology, College of Dentistry, The Ohio State University, Columbus, OH, USA

<sup>2</sup>Laboratory of Clinical Immunology and Microbiology, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Rockville, MD, USA

<sup>3</sup>James Cancer Center, The Ohio State University, Columbus, OH, USA

A supplemental appendix to this article is available online.

## Corresponding Author:

P.S. Kumar, Division of Periodontology, College of Dentistry, The Ohio State University, 4111 Postle Hall, 305, W 12th Avenue, Columbus, OH 43210, USA.

Email: [kumar.83@osu.edu](mailto:kumar.83@osu.edu)

enabled us to fill a critical gap in our knowledge and provide urgently needed information for appropriately apportioning our already stretched resources and guiding best practices for infection control during the COVID-19 pandemic and beyond.

## Methods

### Ethics Statement

This study was approved by the institutional review board of The Ohio State University (protocol 2020H0155) and carried out according to the approved guidelines and in accordance with the STROBE guidelines (Strengthening the Reporting of Observational Studies in Epidemiology) for human observational investigations.

### Participant Selection and Recruitment

Informed consent was obtained from 28 individuals seeking dental treatment in the College of Dentistry of The Ohio State University between May 4 and July 10, 2020. Exclusion criteria were age <18 y, current pregnancy, requirement for antibiotics prior to dental therapy, self-reported HIV or COVID-19 history, COVID-19–like symptoms since January 2020, and antibiotic therapy within 3 mo of sample collection. The sample size was based on previous evidence from in vitro and clinical studies on aerosol spread (King et al. 1997; Muzzini et al. 1999; Nulty et al. 2020; Allison et al. 2021) during scaling and restorative procedures.

### Operating Conditions

The AGDPs were dental implants, restorative procedures with high-speed handpieces, and ultrasonic scaling and were delivered by 5 operators and assistants in 2 enclosed operatories measuring 10.5 × 10 × 12 ft, with 6-exchange/min ventilation. Since aerosols are generated during normal physiologic activity, the operator, assistant, and sample collector wore N95 masks during the entire period to reduce microorganisms from extraneous sources. High-volume intraoral evacuators (mean suction capacity, 7.1 L/min; range, 6.6 to 7.4 L/min) were used throughout the AGDPs. Magnetostrictive ultrasonic scalers were used at a water flow of 19.3 mL/min. Implant osteotomies were performed at a flow rate of 30 mL/min. A high-speed handpiece with a single coolant port and flow rate of 23 mL/min was used for restorative procedures. Twenty-three individuals used a preprocedural rinse of 1% hydrogen peroxide (30 mL for 1 min), and 5 who underwent scaling did not use the rinse.

### Sample Collection

Prior to the procedure, 1 mL of unstimulated saliva and irrigant from the dental (or implant) unit was collected in tubes containing RNA stabilizer (RNALater; Oragene). At 30 min following the procedure, condensate was collected from the face

shields of the operator and assistant, the patient's chest, and an area 6 ft distant from the site of operation ("environment").

### Virus Identification


Investigators conducting the analyses were blinded to the source of samples. An RNA extraction–free dual-plexed reverse transcriptase quantitative polymerase chain reaction method for SARS-CoV-2 detection (SalivaDirect version 5) was used according to the developers' instructions (Vogels et al. 2020). Briefly, 50  $\mu$ L of homogenized saliva was mixed with 2  $\mu$ L of 50-mg/mL proteinase K, and 5  $\mu$ L was used in a 20- $\mu$ L reaction containing FAM-labeled primers and probes targeting SARS-CoV-2 N1 and N2 and amplified for 44 cycles in triplicate reactions. RNA from TRIzol-inactivated virus (obtained from Dr. Wang of The Ohio State University) was used as positive control and to generate standard curves.

### Sequencing and Analysis Pipeline

DNA was isolated from 100  $\mu$ L of the sample with a Qiagen MiniAmp Kit, after a 90-min incubation with lysozyme (2 mg/mL; Thermo Fisher Scientific). V1 to V3 and V4 to V5 (Kumar et al. 2011) regions of the 16S rDNA were sequenced on the Miseq system (2 × 250 bp). Negative and positive controls (defined culture mixture) were used in all runs. Amplicon sequence variants (ASVs) were inferred with the DADA2 version 1.16 pipeline (Callahan et al. 2016). Sequences were truncated on the basis of quality plots and filtered, dereplicated, and denoised with standard parameters, following which chimeras were identified and removed. Paired ends of denoised sequences were then merged. To be retained in the data set, the sequence had to be detected at least once in at least 5% of the samples. ASVs were assigned taxonomic identity with naive Bayes classifiers (QIIME2 q2-feature-classifier; Caporaso et al. 2010) trained for each primer pair by extracting the corresponding hypervariable region from the SILVA database (as of November 12, 2020; Quast et al. 2013).

### Controls

We used multiple internal controls in experimental design as well as analysis to validate these results. As a first level of control, we chose the same irrigant (dental unit water line) that passed through 2 different devices operating on 2 different principles (high-speed handpiece and ultrasonic scaler). We then selected a second therapeutic option, where a different irrigant source (saline) flows through a similar type of device (implant handpiece). By doing so, we controlled for the type of device as well as the type of coolant. In an effort to separate environmental bacteria found normally in water and aerosols from reagent contaminants, we subtracted the sequences found in our negative controls from samples. The negative controls were sterile Petri dishes opened by the sample collector in the clinic environment before the patient, operator, or assistant entered it and subjected to DNA isolation and polymerase



**Figure 1.** Differences in beta diversity between saliva and aerosol deposited on the operator, assistant, patient and the clinic environment. Linear discriminant analysis of Bray-Curtis dissimilarity distances are shown. Two hypervariable regions of the 16S rRNA gene (V1 to V3 and V4 to V5) were sequenced and analyzed separately. Amplicon sequence variants (ASVs) generated from V1-V3 (**A-C**), and V4-V5 regions (**D-F**) are shown. Samples were acquired during (A, D) implant surgery, (B, E) ultrasonic scaling, and (C, F) restorative procedures. Saliva demonstrated significant clustering from all aerosol samples irrespective of procedure ( $P < 0.05$ , Dunn test for joint ranking).

chain reaction amplification in the same manner as the samples. It is well established that the hypervariable region that is sequenced can be a source of bias in determining the composition of the microbiome (Kumar et al. 2011). To overcome this, we used 2 primer pairs targeting 2 hypervariable regions and inducted them into separate sequence analysis pipelines.

### Statistical Analysis


ASVs were used to compute alpha diversity (within group) and beta diversity (between group). As an initial step, the negative controls were used to rarefy each data set. However, the sequencing depth still varied among the samples by a magnitude of 100 in some cases, especially between saliva and environmental condensate. Since current evidence does not support rarefying the microbiome to compensate for sequencing effort (McMurdie and Holmes 2014), we used cumulative sum scaling normalization from the Bioconductor package metagenomeSeq. Linear discriminant analysis (LDA) was performed with the MASS package for R. The input for LDA was a matrix of variance-stabilized (arc-sin square root transformed) relative abundances of ASVs (Shchipkova et al. 2010). MASS:lda provided singular value decomposition values, which were used to calculate the percentage variance explained in each dimension. The Dunn test with joint ranking was used to test significance of LDA clustering. SourceTracker version 0.9.5 (Knights et al. 2011) was applied to identify the source of microorganisms. Saliva and irrigant were designated as possible sources, and the operator, assistant, patient, and ambient

environment were set as targets. Data sets were filtered to remove ASVs that were not present in at least 1% of samples. The following default parameters were used for analysis: rarefaction depth, 1,000; burn-in, 100; restart, 10; and alpha (0.001) and beta (0.01) Dirichlet hyperparameters.

### Results

A total of 4,500,063 classifiable sequences representing 22,013 ASVs were used for microbial analysis. We began by characterizing the microbiota in saliva and the aerosol deposited on the operator, assistant, patient, and environment with an increasingly granular top-down approach. We found significant group separation between the salivary and aerosol microbiomes irrespective of the type of aerosol-generating procedure ( $P < 0.001$ , Dunn method for joint ranking of Bray-Curtis dissimilarity distances; Fig. 1, Appendix Table 1). By contrast, no such clustering was demonstrable among the aerosol deposited on the operator, assistant, patient, and environment. We then explored the abundances of individual taxa in the condensate generated from each procedure. Irrespective of the procedure or the area of deposition, 70% of the microbial abundance was attributable to 2 ASVs that mapped to uncultured *Vulcaniibacteria*. These ASVs were also the predominant taxa in the irrigants. Salivary bacteria, when present, accounted for 0.1% to 1.2% of the microbial abundance of aerosol.

Source-tracking analysis revealed that, irrespective of the AGDP, microbiota from irrigants contributed to a median 78% of the microbiota in condensate (range, 2.5% to 100%; Fig. 2),



**Figure 2.** Source of microorganisms in aerosols generated during dental procedures. The relative contributions of saliva, irrigant fluid and unknown sources to the microbial composition of the condensate deposited on the assistant (A), environment (B), operator (C), and patient (D) are shown. The proportions were estimated using SourceTracker. For each subject, irrigant and saliva were designated as potential sources, and the operator, assistant, patient and the ambient environment were set as targets. Datasets were filtered to remove taxa that were not present in at least 1% of samples. Default parameters (rarefaction depth 1000, burn-in 100, restart 10, alpha [0.001] and beta [0.01] dirichlet hyperparameter) were used for analysis.

while saliva contributed to a median 0% (range, 0% to 82%). On average, 20% of the microbiota could not be attributed to either source (range, 0% to 90%). Salivary bacteria were detectable in the condensate in only 8 participants out of the 28. Of these, 5 had not used a preprocedural rinse. In these 8 cases, the patient's chest was the most frequent site of deposition ( $P < 0.05$ , chi-square test). LDA and SourceTracker revealed similar trends for ASVs generated from V1 to V3 and V4 to V5, indicating that our findings were robust against sequencing targets (Fig. 1).

We then investigated if aerosol dispersion of SARS-CoV-2 virus followed the same pattern, and it was identified in the saliva of 19 participants, with a viral load between 27 and 912 copies/mL. The virus was undetectable in the condensate on the operator, assistant, patient, or environment in any of these cases (Table).

## Discussion

The impetus for this study came from our recent review of the literature on aerosols generated during dental procedures (Kumar and Subramanian 2020). We found that while several investigations measured the radius of contamination following AGDPs and demonstrated the presence of microbiota in these contaminated areas (Grenier 1995; Harrel et al. 1998; Rivera-Hidalgo et al. 1999; Petti and Vitali 2017; Kobza et al. 2018; Jain et al. 2020), few characterized the types of microorganisms, and none identified their source. Understanding the sources of microbial bioload in aerosols is of immediate urgency, not only for infection control in dental operatories during the COVID-19 pandemic, but also to inform best practices in aerosol reduction, mitigation, and abatement in the long term. To the best of our knowledge, this study sets the stage for future work on risk of disease transmission among dental health care workers and patients.

We deliberately elected to employ a passive air-sampling methodology over other aerosol-harvesting methods (Greco and Lai 2008; Verreault et al. 2008; O'Neil et al. 2017), since dental procedures generate a variety of airborne particles, ranging from spatter and particulate matter to aerosolized microorganisms, any of which could pose a health hazard to patient and personnel. Our method allowed us to collect everything that was generated during the AGDP as well as everything that settled on the surfaces during the following 30 min. This method has been widely used in similar studies, since it is reflective of events that occur in a treatment session (Bentley et al. 1994; Bennett et al. 2000; Hallier et al. 2010).

This study was conducted at the height of the COVID-19 pandemic, and patients were inducted into the study based on presentation for treatment to the dental clinics. We did not do a COVID-19 test before starting the study because the primers and probes had not been optimized when we began our recruitment. An extensive review of the literature revealed wide heterogeneity in sample size, with a minimum of 3 (Bentley et al. 1994) and a maximum of 20 (Serban et al. 2013; Singh et al. 2016). As a result, we included 15 participants for the ultrasonic group. During the same time frame, 10 patients who presented for implant therapy consented to participate. Thus, we did an interim analysis to see if we could detect differences among the aerosols deposited on the various surfaces with 10 participants, and since there were significant differences, we used this as our endpoint. Three studies used 3 subjects/replicates per group for restorative procedures and to estimate SARS-CoV-2 spread (van Doremalen et al. 2020; Allison et al. 2021) and therefore we included the restorative data from 3 subjects for completeness.

With all the procedural controls described in the methods, we used an emergent bioinformatics approach to characterize microbial assemblages: ASVs. Bioinformatics has traditionally sought to reduce large 16S sequence data sets into manageable

**Table.** Levels of SARS-CoV-2 Virus in 28 Asymptomatic Patients with Noncontributory COVID-19 History and in Aerosols Generated during Dental Procedures on Them.

| ID               | Procedure  | Periodontal Status                 | Saliva     |            | Operator   |            | Assistant  |            | Environment |            | Patient    |            |
|------------------|------------|------------------------------------|------------|------------|------------|------------|------------|------------|-------------|------------|------------|------------|
|                  |            |                                    | Average Ct | Viral Load | Average Ct | Viral Load | Average Ct | Viral Load | Average Ct  | Viral Load | Average Ct | Viral Load |
| S1               | Ultrasonic | Generalized moderate periodontitis | 40.00      | ND         | NA         | NA         | NA         | NA         | NA          | NA         | NA         | NA         |
| S10 <sup>a</sup> | Ultrasonic | Periodontal health                 | 36.61      | 109.4      | 40         | ND         | 40         | ND         | 40          | ND         | 40         | ND         |
| S11              | Ultrasonic | Periodontal health                 | 37.41      | 62.2       | 40         | ND         | 40         | ND         | 40          | ND         | 40         | ND         |
| S12              | Drilling   | Periodontal health                 | 38.45      | 29.9       | 40         | ND         | 40         | ND         | 40          | ND         | 40         | ND         |
| S13              | Ultrasonic | Periodontal health                 | 37.01      | 82.2       | 40         | ND         | 40         | ND         | 40          | ND         | 40         | ND         |
| S14              | Ultrasonic | Generalized moderate periodontitis | 35.01      | 338.2      | 40         | ND         | 40         | ND         | 40          | ND         | 40         | ND         |
| S16              | Implant    | Periodontal health                 | 40.00      | ND         | NA         | NA         | NA         | NA         | NA          | NA         | NA         | NA         |
| S17              | Implant    | Periodontal health                 | 40.00      | ND         | NA         | NA         | NA         | NA         | NA          | NA         | NA         | NA         |
| S18              | Ultrasonic | Periodontal health                 | 37.12      | 76.3       | 40         | ND         | 40         | ND         | 40          | ND         | 40         | ND         |
| S19              | Ultrasonic | Generalized moderate periodontitis | 37.14      | 75.4       | 40         | ND         | 40         | ND         | 40          | ND         | 40         | ND         |
| S2               | Implant    | Periodontal health                 | 40.00      | ND         | NA         | NA         | NA         | NA         | NA          | NA         | NA         | NA         |
| S20              | Ultrasonic | Generalized moderate periodontitis | 35.81      | 192.6      | 40         | ND         | 40         | ND         | 40          | ND         | 40         | ND         |
| S21 <sup>a</sup> | Ultrasonic | Generalized moderate periodontitis | 36.30      | 136.1      | 40         | ND         | 40         | ND         | 40          | ND         | 40         | ND         |
| S22 <sup>a</sup> | Ultrasonic | Generalized moderate periodontitis | 33.60      | 912.5      | 40         | ND         | 40         | ND         | 40          | ND         | 40         | ND         |
| S23              | Implant    | Periodontal health                 | 36.07      | 160.4      | 40         | ND         | 40         | ND         | 40          | ND         | 40         | ND         |
| S24              | Ultrasonic | Periodontal health                 | 37.27      | 68.7       | 40         | ND         | 40         | ND         | 40          | ND         | 40         | ND         |
| S26 <sup>a</sup> | Ultrasonic | Generalized moderate periodontitis | 35.09      | 318.6      | 40         | ND         | 40         | ND         | 40          | ND         | 40         | ND         |
| S27 <sup>a</sup> | Ultrasonic | Periodontal health                 | 36.92      | 88.0       | 40         | ND         | 40         | ND         | 40          | ND         | 40         | ND         |
| S28              | Drilling   | Periodontal health                 | 37.83      | 46.2       | 40         | ND         | 40         | ND         | 40          | ND         | 40         | ND         |
| S29              | Implant    | Periodontal health                 | 40.00      | ND         | NA         | NA         | NA         | NA         | NA          | NA         | NA         | NA         |
| S3               | Implant    | Periodontal health                 | 37.02      | 81.9       | 40         | ND         | 40         | ND         | 40          | ND         | 40         | ND         |
| S30              | Implant    | Periodontal health                 | 38.56      | 27.7       | 40         | ND         | 40         | ND         | 40          | ND         | 40         | ND         |
| S4               | Implant    | Periodontal health                 | 40.00      | ND         | NA         | NA         | NA         | NA         | NA          | NA         | NA         | NA         |
| S5               | Implant    | Periodontal health                 | 40.00      | ND         | NA         | NA         | NA         | NA         | NA          | NA         | NA         | NA         |
| S6               | Implant    | Periodontal health                 | 40.00      | ND         | NA         | NA         | NA         | NA         | NA          | NA         | NA         | NA         |
| S7               | Ultrasonic | Periodontal health                 | 40.00      | ND         | NA         | NA         | NA         | NA         | NA          | NA         | NA         | NA         |
| S8               | Implant    | Periodontal health                 | 36.57      | 112.3      | 40         | ND         | 40         | ND         | 40          | ND         | 40         | ND         |
| S9               | Drilling   | Periodontal health                 | 37.53      | 57.2       | 40         | ND         | 40         | ND         | 40          | ND         | 40         | ND         |

NA, aerosol sample not analyzed since the virus was not identified in the saliva samples; ND, not detected by reverse transcriptase quantitative polymerase chain reaction at a detection threshold of 40 reaction cycles.

<sup>a</sup>Patient did not use a mouth rinse.

operational units, either by assigning them a taxonomic identity by comparison with a database or by identifying taxon-independent operational taxonomic units with a sequence similarity threshold (Huse et al. 2010). Both these approaches fail to capture the immense diversity of environmental bacteria, many of which have not been assigned definitive taxonomic identities or entered into curated databases (Callahan et al. 2017). The decision to use ASVs over operational taxonomic units was driven by the need for a finer scale resolution of bacterial identities than could be provided by traditional methods. This allowed us to exploit the immense sensitivity

built into the 16S rRNA gene and overcome the limitations imposed by incomplete reference databases to better discriminate ecologic patterns (Prodan et al. 2020).

Our key discovery was that the irrigant fluid contributes to the majority of the bioload in dental aerosols. Although this is the first time that it is being reported, this is not entirely surprising, since we estimated that the irrigant flow dilutes the saliva by 20- to 200-fold (assuming a salivary flow rate of 0.1 to 1.0 mL/min; Iorgulescu 2009). This discovery lends credence to previous studies on the transmission of legionellosis, a waterborne infection, and pneumonia, from dental unit water

lines (Ricci et al. 2012; Petti and Vitali 2017). It also validates recent reports of extremely low COVID-19 transmission among dental personnel (Estrich et al. 2020).

One surprising finding was the discovery of microorganisms in the implant irrigant, which was sterile saline. To confirm that this was not an artifact, we sequenced the saline before and after it passed through the implant handpiece and discovered a diverse microbiome in reusable handpieces. However, since these handpieces were sterilized by autoclaving, we anticipate that these organisms are not viable.

The salivary SARS-CoV-2 load in our study is consistent with previous reports of  $10^2$  to  $10^6$  copies/mL (Chau et al. 2020); however, none was detected in any aerosol. Interestingly, participant 22 demonstrated the highest salivary viral load, and while bacteria of salivary origin could be identified on the operator and environment, the virus was not detectable in the same samples. This was true of those individuals with periodontitis and those who were periodontally healthy. The 20- to 200-fold dilution factor and our discovery that the irrigant, not the saliva, is the primary and predominant source of airborne microbiota might explain the absence of SARS-CoV-2 in aerosol. Preprocedural mouth rinsing serves to further reduce potential salivary contaminants in aerosol.

In summary, we find sufficient evidence to reject our null hypothesis that saliva is a potential source of disease transmission during AGDPs. Instead, we identify, for the first time, that the irrigant used in dental equipment is the primary and major source of microorganisms during AGDPs. We also demonstrate that high-volume intraoral evacuators are highly effective in reducing salivary contamination of the surrounding environment and that preprocedural mouth rinses consistently reduce salivary microbial bioloads. Within the limitations of a small sample size, we conclude that the risk for transmission of SARS-CoV-2 and other respiratory pathogens from aerosolized saliva in dental operatories is moderately low and that current infection control practices are adequately robust to protect personnel and patients alike.

### Author Contributions

A.P. Meethil, S. Saraswat, P.P. Chaudhary, S.M. Dabdoub, P.S. Kumar, contributed to conception, design, and data acquisition, drafted and critically revised manuscript. All authors gave final approval and agree to be accountable for all aspects of the work.

### Declaration of Conflicting Interests


The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

### Funding

The authors received no financial support for the research, authorship, and/or publication of this article.

### ORCID iDs

S.M. Dabdoub  <https://orcid.org/0000-0002-1374-8347>

P.S. Kumar  <https://orcid.org/0000-0001-5844-1341>

### References

- Allison JR, Currie CC, Edwards DC, Bowes C, Coulter J, Pickering K, Kozhevnikova E, Durham J, Nile CJ, Jakubovics N, et al. 2021. Evaluating aerosol and splatter following dental procedures: addressing new challenges for oral health care and rehabilitation. *J Oral Rehabil*. 48(1):61–72.
- Bennett AM, Fulford MR, Walker JT, Bradshaw DJ, Martin MV, Marsh PD. 2000. Microbial aerosols in general dental practice. *Br Dent J*. 189(12):664–667.
- Bentley CD, Burkhart NW, Crawford JJ. 1994. Evaluating spatter and aerosol contamination during dental procedures. *J Am Dent Assoc*. 125(5):579–584.
- Callahan BJ, McMurdie PJ, Holmes SP. 2017. Exact sequence variants should replace operational taxonomic units in marker-gene data analysis. *ISME J*. 11(12):2639–2643.
- Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. 2016. DADA2: high-resolution sample inference from illumina amplicon data. *Nat Methods*. 13(7):581–583.
- Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, Fierer N, Pena AG, Goodrich JK, Gordon JL, et al. 2010. QIIME allows analysis of high-throughput community sequencing data. *Nat Methods*. 7(5):335–336.
- Centers for Disease Control and Prevention. 2020. Interim infection prevention and control guidance for dental settings during the COVID-19 response [accessed 2020 Jul 14]. <https://www.cdc.gov/coronavirus/2019-ncov/hcp/dental-settings.html>
- Chau NVV, Thanh Lam V, Thanh Dung N, Yen LM, Minh NNQ, Hung LM, Ngoc NM, Dung NT, Man DNH, Nguyet LA, et al.; Oxford University Clinical Research Unit COVID-19 Research Group. 2020. The natural history and transmission potential of asymptomatic SARS-CoV-2 infection. *Clin Infect Dis*. 71(10):2679–2687.
- D'Aiuto F, Graziani F, Tete S, Gabriele M, Tonetti MS. 2005. Periodontitis: from local infection to systemic diseases. *Int J Immunopathol Pharmacol*. 18(3 Suppl):1–11.
- Davies A, Thomson G, Walker J, Bennett A. 2009. A review of the risks and disease transmission associated with aerosol generating medical procedures. *J Infect Prev*. 10(4):122–126.
- de Pablo P, Chapple IL, Buckley CD, Dietrich T. 2009. Periodontitis in systemic rheumatic diseases. *Nat Rev Rheumatol*. 5(4):218–224.
- Estrich CG, Mikkelsen M, Morrissey R, Geisinger ML, Ioannidou E, Vujicic M, Araujo MWB. 2020. Estimating COVID-19 prevalence and infection control practices among US dentists. *J Am Dent Assoc*. 151(11):815–824.
- Greco PM, Lai C-H. 2008. A new method of assessing aerosolized bacteria generated during orthodontic debonding procedures. *Am J Orthod Dentofacial Orthop*. 133(4):S79–S87.
- Grenier D. 1995. Quantitative analysis of bacterial aerosols in two different dental clinic environments. *Appl Environ Microbiol*. 61(8):3165–3168.
- Gupta S, Sahni V. 2020. The intriguing commonality of NETosis between COVID-19 & periodontal disease. *Med Hypotheses*. 144:109968.
- Hallier C, Williams DW, Potts AJC, Lewis MAO. 2010. A pilot study of bio-aerosol reduction using an air cleaning system during dental procedures. *Br Dent J*. 209(8):E14.
- Harrel SK, Barnes JB, Rivera-Hidalgo F. 1998. Aerosol and splatter contamination from the operative site during ultrasonic scaling. *J Am Dent Assoc*. 129(9):1241–1249.
- Huse SM, Welch DM, Morrison HG, Sogin ML. 2010. Ironing out the wrinkles in the rare biosphere through improved OTU clustering. *Environ Microbiol*. 12(7):1889–1898.
- Iorgulescu G. 2009. Saliva between normal and pathological: important factors in determining systemic and oral health. *J Med Life*. 2(3):303–307.
- Jain M, Mathur A, Mathur A, Mukhi PU, Ahire M, Pingal C. 2020. Qualitative and quantitative analysis of bacterial aerosols in dental clinical settings: risk exposure towards dentist, auxiliary staff, and patients. *J Family Med Prim Care*. 9(2):1003–1008.
- Judson SD, Munster VJ. 2019. Nosocomial transmission of emerging viruses via aerosol-generating medical procedures. *Viruses*. 11(10):940.
- Kassebaum NJ, Bernabe E, Dahiya M, Bhandari B, Murray CJ, Marcenes W. 2014. Global burden of severe periodontitis in 1990–2010: a systematic review and meta-regression. *J Dent Res*. 93(11):1045–1053.
- King TB, Muzzin KB, Berry CW, Anders LM. 1997. The effectiveness of an aerosol reduction device for ultrasonic scalers. *J Periodontol*. 68(1):45–49.
- Knights D, Kuczynski J, Charlson ES, Zaneveld J, Mozer MC, Collman RG, Bushman FD, Knight R, Kelley ST. 2011. Bayesian community-wide culture-independent microbial source tracking. *Nat Methods*. 8(9):761–763.
- Kobza J, Pastuszka JS, Bragoszewska E. 2018. Do exposures to aerosols pose a risk to dental professionals? *Occup Med (Lond)*. 68(7):454–458.
- Kumar PS, Brooker MR, Dowd SE, Camerlengo T. 2011. Target region selection is a critical determinant of community fingerprints generated by 16S pyrosequencing. *PLoS One*. 6(6):e20956.



- Kumar PS, Subramanian K. 2020. Demystifying the mist: sources of microbial bioload in dental aerosols. *J Periodontol.* 91(9):1113–1122.
- McMurdie PJ, Holmes S. 2014. Waste not, want not: why rarefying microbiome data is inadmissible. *PLoS Comput Biol.* 10(4):e1003531.
- Muzzin KB, King TB, Berry CW. 1999. Assessing the clinical effectiveness of an aerosol reduction device for the air polisher. *J Am Dent Assoc.* 130(9):1354–1359.
- National Institute of Dental and Craniofacial Research. 2014. Tooth loss among adults 20–64 [accessed 2020 Dec 17]. <http://www.nidcr.nih.gov/DataStatistics/FindDataByTopic/ToothLoss/ToothLossAdults20to64.htm>
- Nulty A, Lefkadtis C, Zachrisson P, Van Tonder Q, Yar R. 2020. A clinical study measuring dental aerosols with and without a high-volume extraction device. *Br Dent J* [epub ahead of print 20 Nov 2020]. doi:10.1038/s41415-020-2274-3
- O’Neil CA, Li J, Leavey A, Wang Y, Hink M, Wallace M, Biswas P, Burnham C-AD, Babcock HM, et al.; Centers for Disease Control and Prevention. 2017. Characterization of aerosols generated during patient care activities. *Clin Infect Dis.* 65(8):1335–1341.
- Petti S, Vitali M. 2017. Occupational risk for legionella infection among dental healthcare workers: meta-analysis in occupational epidemiology. *BMJ Open.* 7(7):e015374.
- Pitones-Rubio V, Chávez-Cortez EG, Hurtado-Camarena A, González-Rascón A, Serafin-Higuera N. 2020. Is periodontal disease a risk factor for severe COVID-19 illness? *Med Hypotheses.* 144:109969.
- Prodan A, Tremaroli V, Brolin H, Zwinderman AH, Nieuwdorp M, Levin E. 2020. Comparing bioinformatic pipelines for microbial 16S rRNA amplicon sequencing. *PLoS One.* 15(1):e0227434.
- Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glockner FO. 2013. The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Res.* 41(Database Issue):D590–D596.
- Ricci ML, Fontana S, Pinci F, Fiumana E, Pedna MF, Farolfi P, Sabattini MA, Scaturro M. 2012. Pneumonia associated with a dental unit waterline. *Lancet (Lond, Engl).* 379(9816):684.
- Rivera-Hidalgo F, Barnes JB, Harrel SK. 1999. Aerosol and splatter production by focused spray and standard ultrasonic inserts. *J Periodontol.* 70(5):473–477.
- Serban D, Banu A, Serban C, Tuta-Sas I, Vlaicu B. 2013. Predictors of quantitative microbiological analysis of spatter and aerosolization during scaling. *Rev Med Chir Soc Med Nat Iasi.* 117(2):503–508.
- Shchipkova AY, Nagaraja HN, Kumar PS. 2010. Subgingival microbial profiles in smokers with periodontitis. *J Dent Res.* 89(11):1247–1253.
- Singh A, Shiva Manjunath RG, Singla D, Bhattacharya HS, Sarkar A, Chandra N. 2016. Aerosol, a health hazard during ultrasonic scaling: a clinico-microbiological study. *Indian J Dent Res.* 27(2):160–162.
- Tonetti MS, Van Dyke TE; Working Group 1 of the Joint EFP/AAP Workshop. 2013. Periodontitis and atherosclerotic cardiovascular disease: consensus report of the Joint EFP/AAP Workshop on Periodontitis and Systemic Diseases. *J Periodontol.* 84(4):S24–S29.
- van Doremalen N, Bushmaker T, Morris DH, Holbrook MG, Gamble A, Williamson BN, Tamin A, Harcourt JL, Thornburg NJ, Gerber SI, et al. 2020. Aerosol and surface stability of SARS-CoV-2 as compared with SARS-CoV-1. *N Engl J Med.* 382(16):1564–1567.
- Verreault D, Moineau S, Duchaine C. 2008. Methods for sampling of airborne viruses. *Microbiol Mol Biol Rev.* 72(3):413–444.
- Vogels C, Brackney DE, Kalinich C, Ott I, Grubaugh N, Wyllie A. 2020. Salivadirect: RNA extraction-free SARS-CoV-2 diagnostics [accessed 2021 Apr 20]. <https://www.protocols.io/view/salivadirect-ma-extraction-free-sars-cov-2-diagno-btdnni5e>
- Zhao S, Cao J, Sun R, Zhang L, Liu B. 2020. Analysis of anxiety-related factors amongst frontline dental staff during the COVID-19 pandemic in Yichang, China. *BMC Oral Health.* 20(1):342.